

STABILITY OF TWO DOUBLET ELECTROWEAK STRINGS

MICHAEL A. EARNSHAW

MARGARET JAMES

Department of Applied Mathematics and Theoretical Physics

Silver Street, Cambridge, CB3 9EW, Great Britain

ABSTRACT

Vortex solutions in the two Higgs doublet electroweak model are constructed, and their stability to small perturbations is studied. The most general perturbation is decomposed into angular momentum modes, the least stable mode is identified, and the linearised energy change of the vortex under this perturbation is calculated numerically for various choices of parameters, thus determining whether or not the string is stable. It is found that, for realistic values of the Higgs mass and Weinberg angle, the string is unstable.

1. Introduction

In the last two years, interest has been generated in the study of ‘embedded’ defects. These occur when a theory which has topological defects is contained in a larger theory and the defects remain solutions of the field equations. General conditions for this to take place have been studied by Vachaspati et al^[1]. A crucial question is whether the defect is still stable. Generally we will not expect them to remain topologically stable and the stability* becomes a dynamical question.

In this paper we consider embedded vortex solutions. A prototype example is the semi-local string^[2]; the Nielsen-Olesen vortex^[3] is embedded in the semilocal model where the gauge group $SU(2)_{global} \times U(1)_{local}$ is spontaneously broken to $U(1)_{global}$ by a complex Higgs doublet. The vacuum manifold of the Higgs is now S^3 and since the first homotopy group of this is trivial the solution is no longer topologically stable. However the solutions may still be classified into topological sectors separated by infinite energy since finite energy requires that the Higgs field at spatial infinity lies on a $U(1)$ orbit on S^3 . The question is then whether the solution is dynamically stable in a given sector. There is one free parameter in the model, the ratio of the quartic coupling to the gauge coupling and the string is stable for a finite parameter range.^[4,5]

When the $SU(2)$ symmetry is gauged we recover the electroweak model, $SU(2)_L \times U(1)_Y \rightarrow U(1)_{em}$. The vortex solution persists in the model for any value of the Weinberg angle^[6,7]. Now all configurations are in the vacuum sector i.e. any solution can be continuously deformed to the vacuum. The parameter space is now two-dimensional, the parameters being the ratio of the Higgs-mass to the Z-mass and the Weinberg angle. The string has been shown to be stable for a range in parameter space^[8,9]. However taking the experimental lower bound on the scalar Higgs and the observed value of the Weinberg angle, it was shown that if the minimal standard model is the physically realised model then electroweak-strings are unstable. The calculation was refined by including finite temperature and quantum effects^[10] and while the stability

* We are only considering the stability under small perturbations. Hence, the word ‘stability’ should always be understood as ‘meta-stability’.

of the string is improved the physical values are still outside the region of stability for temperatures close to the electroweak phase transition. This is somewhat disappointing since the prospect of a stable classical vortex is an exciting one. It could possibly be produced in future particle accelerators. However it is not yet understood how to calculate the cross-section for what, in quantum theory would be a coherent state of many particles and it may well be that the available phase space is so small as to make the cross-section negligible. In cosmology if the strings are meta-stable, there would be implications for physics at the electroweak phase transition. Davis and Brandenberger^[11] have suggested that they could play a role in baryogenesis. We are therefore led to consider extensions of the standard model where the stability region might be enlarged sufficiently to include the physical values. Vachaspati and Watkins^[12] suggested that the vortex could be stabilised by the introduction of a further scalar field which would form a bound state with the string.

A different possibility, the one we investigate here, is that of extending the standard model via the introduction of a second Higgs doublet. For simplicity we do not consider the most general such model but restrict ourselves by imposing CP-invariance on the Higgs potential. The model has seven parameters: the Weinberg angle, five Higgs couplings and the ratio of the norm of the vacuum expectation values of the two Higgs doublets. The stability analysis follows the same scheme as that in reference [8] although the introduction of the second Higgs doublet somewhat complicates the analysis. In section 2. we discuss the vortex solutions in the two Higgs model. In section 3. we consider the perturbations around the string solution and show that they decompose into channels labelled by angular momentum and the eigenvalues of two discrete symmetry operators. In section 4. we locate the least stable mode and reduce the analysis to solving a pair of coupled Schrödinger equations. In section 5. we discuss the numerical work and present our results and in 6. we give our conclusions.

2. Two Higgs-Doublet Electroweak Strings

We consider the minimal extension of the Higgs sector of the standard model by introducing a second Higgs complex doublet. We give the most general potential subject only to the restrictions of gauge invariance and that it has the discrete symmetry, $(\Phi_1 \leftrightarrow -\Phi_1, \Phi_2 \leftrightarrow -\Phi_2)$ ^[13]. (The latter is required to ensure that flavour changing neutral currents are not too large.)

$$\begin{aligned} V(\Phi_1, \Phi_2) = & \frac{1}{2}\lambda_1(|\Phi_1|^2 - v_1^2)^2 + \frac{1}{2}\lambda_2(|\Phi_2|^2 - v_2^2)^2 + \frac{1}{2}\lambda_3(|\Phi_1|^2 + |\Phi_2|^2 - v_1^2 - v_2^2)^2 \\ & + \lambda_4 \left((\Phi_1^\dagger \Phi_1)(\Phi_2^\dagger \Phi_2) - |\Phi_1^\dagger \Phi_2|^2 \right) + \lambda_5 \left(\text{Re}(\Phi_1^\dagger \Phi_2) - v_1 v_2 \cos \eta \right)^2 \\ & + \lambda_6 \left(\text{Im}(\Phi_1^\dagger \Phi_2) - v_1 v_2 \sin \eta \right)^2 \end{aligned} \quad (1)$$

The vacuum expectation of the Higgs fields are given by

$$\langle \Phi_1 \rangle = \begin{pmatrix} 0 \\ v_1 \end{pmatrix}, \quad \langle \Phi_2 \rangle = \begin{pmatrix} 0 \\ v_2 e^{i\eta} \end{pmatrix}. \quad (2)$$

If $\sin \eta \neq 0$ then there is CP-violation in the Higgs sector. This case is considerably more complicated than the $\sin \eta = 0$ case where, as we shall see later, the vortex solution is CP-invariant and the perturbations then split into channels labelled CP-even and CP-odd. Hence we set $\sin \eta = 0$. Also, the λ_6 term is the mass term for the CP-odd neutral scalar, A_0 , which will decouple from our analysis, so we are free to set $\lambda_6 = \lambda_5$. Then the last two terms are replaced by a single term

$$\lambda_5 \left| (\Phi_1^\dagger \Phi_2) - v_1 v_2 \right|^2. \quad (3)$$

We are considering static, bosonic field configurations. There is no time dependence and we choose a gauge where the zero components of the gauge fields are set to zero. The energy functional is given by

$$E = \int d^3x \left[\frac{1}{4} G_{ij}^a G_{ij}^a + \frac{1}{4} F_{ij} F_{ij} + (D_j \Phi_{\bar{a}})^\dagger (D_j \Phi_{\bar{a}}) + V(\Phi_1, \Phi_2) \right] \quad (4)$$

where $\bar{a} = 1, 2^*$ and

$$G_{ij}^a = \partial_i W_j^a - \partial_j W_i^a + g \epsilon^{abc} W_i^b W_j^c, \quad (5)$$

* Barred indices always take the values 1 and 2.

$$F_{ij} = \partial_i B_j - \partial_j B_i, \quad (6)$$

$$D_i \Phi_{\bar{a}} = \partial_i \Phi_{\bar{a}} - \frac{1}{2} i g \tau^a W_i^a \Phi_{\bar{a}} - \frac{1}{2} i g' B_i \Phi_{\bar{a}} \quad (7)$$

where we take $T^a = -\frac{1}{2} i \tau^a$ as our basis for $L(SU(2))$ (denoting the Lie algebra of $SU(2)$) with $[T^a, T^b] = \epsilon^{abc} T^c$. The Weinberg angle is given by $\tan \Theta_W = g'/g$. The semiclassical masses of the W-boson and the Z-boson are, respectively,

$$M_W^2 = \frac{1}{2} g^2 (v_1^2 + v_2^2), \quad M_Z^2 = \frac{1}{2} \alpha^2 (v_1^2 + v_2^2), \quad (8)$$

where $\alpha = (g'^2 + g^2)^{\frac{1}{2}}$.

There are five physical Higgs bosons, a charged pair $H^\pm$, two CP-even scalars, H^0 and h^0 , and the CP-odd neutral scalar A_0 which we have already discussed. The charged pair have mass $m_{H^\pm}^2 = \lambda_4 (v_1^2 + v_2^2)$ and the two remaining Higgs scalars mix through the mass matrix,

$$M = \begin{pmatrix} 2v_1^2(\lambda_1 + \lambda_3) + v_2^2\lambda_5 & (2\lambda_3 + \lambda_5)v_1v_2 \\ (2\lambda_3 + \lambda_5)v_1v_2 & 2v_2^2(\lambda_2 + \lambda_3) + v_1^2\lambda_5 \end{pmatrix} \quad (9)$$

and the physical eigenstates have masses

$$m_{H^0, h^0}^2 = \frac{1}{2} (M_{11} + M_{22} \pm ((M_{11} - M_{22})^2 + 4M_{12}^2)^{1/2}) \quad (10)$$

where by convention we take $m_{H^0} > m_{h^0}$.

The time-independent field equations are

$$\begin{aligned} D_j G_{ij}^a &= -\frac{1}{2} i g \left(\Phi_{\bar{a}}^\dagger \tau^a D_i \Phi_{\bar{a}} - (D_i \Phi_{\bar{a}})^\dagger \tau^a \Phi_{\bar{a}} \right) \\ \partial_j F_{ij} &= -\frac{1}{2} i g' \left(\Phi_{\bar{a}}^\dagger D_i \Phi_{\bar{a}} - (D_i \Phi_{\bar{a}})^\dagger \Phi_{\bar{a}} \right) \\ D_i D_i \Phi_1 &= \lambda_1 (|\Phi_1|^2 - v_1^2) \Phi_1 + \lambda_3 (|\Phi_1|^2 + |\Phi_2|^2 - v_1^2 - v_2^2) \Phi_1 \\ &\quad + \lambda_4 (|\Phi_2|^2 \Phi_1 - (\Phi_2^\dagger \Phi_1) \Phi_2) + \lambda_5 (\Phi_2^\dagger \Phi_1 - v_1 v_2) \Phi_2 \\ D_i D_i \Phi_2 &= \lambda_2 (|\Phi_2|^2 - v_2^2) \Phi_2 + \lambda_3 (|\Phi_1|^2 + |\Phi_2|^2 - v_1^2 - v_2^2) \Phi_2 \\ &\quad + \lambda_4 (|\Phi_1|^2 \Phi_2 - (\Phi_1^\dagger \Phi_2) \Phi_1) + \lambda_5 (\Phi_1^\dagger \Phi_2 - v_1 v_2) \Phi_1 \end{aligned} \quad (11)$$

In addition, we recall the usual mixing formulae:

$$Z_i \equiv \cos \Theta_W W_i^3 - \sin \Theta_W B_i, \quad A_i \equiv \sin \Theta_W W_i^3 + \cos \Theta_W B_i, \quad (12)$$

The vortex solution extremising (4) is given by :

$$\begin{aligned}\vec{W}^1 = 0 = \vec{W}^2 = \vec{A}, \quad \vec{Z} = -\frac{Z(r)}{r} \hat{e}_\theta \\ \Phi_1 = f_1(r) e^{im_1\theta} \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad \Phi_2 = f_2(r) e^{im_2\theta} \begin{pmatrix} 0 \\ 1 \end{pmatrix},\end{aligned}\tag{13}$$

where the coordinates r and θ are polar coordinates in the xy -plane and m_1 and m_2 are integers. On substitution of (13) into the equations of motion they reduce to

$$\begin{aligned}f_1'' + \frac{f_1'}{r} - \left(m_1 - \frac{\alpha}{2}Z\right)^2 \frac{f_1}{r^2} - (\lambda_1 + \lambda_3)(f_1^2 - v_1^2)f_1 \\ - \lambda_3(f_2^2 - v_2^2)f_1 - \lambda_5(f_1f_2 - v_1v_2)f_2 = 0\end{aligned}\tag{14}$$

$$\begin{aligned}f_2'' + \frac{f_2'}{r} - \left(m_2 - \frac{\alpha}{2}Z\right)^2 \frac{f_2}{r^2} - (\lambda_2 + \lambda_3)(f_2^2 - v_2^2)f_2 \\ - \lambda_3(f_1^2 - v_1^2)f_2 - \lambda_5(f_1f_2 - v_1v_2)f_1 = 0\end{aligned}\tag{15}$$

$$Z'' - \frac{Z'}{r} + \alpha \left(\left(m_1 - \frac{\alpha}{2}Z\right) f_1^2 + \left(m_2 - \frac{\alpha}{2}Z\right) f_2^2 \right) = 0\tag{16}$$

where primes denote differentiation with respect to r . It is clear from the equations that we must have $m_1 = m_2$. This can also be seen by examining the energy functional and noting that to prevent a logarithmic divergence in the energy integral at infinity, we require that the angular dependence of the two Higgs doublets must be identical asymptotically. We shall restrict ourselves to the case $m_1 = m_2 = 1$. The equations are then solved together with the boundary conditions:

$$f_1(0) = f_2(0) = Z(0) = 0, \quad \frac{f_1(\infty)}{v_1} = \frac{f_2(\infty)}{v_2} = \frac{\alpha}{2}Z(\infty) = 1\tag{17}$$

Asymptotically, for large r , writing $f_1 = v_1 + \delta_1$, $f_2 = v_2 + \delta_2$ and $Z = \frac{2}{\alpha} + r\delta_3$ and linearising the equations of motion when $m_{H_0}, m_{h_0} < m_Z$, for example when λ_i/α^2 are small, we obtain the following modified Bessel equations

$$\begin{pmatrix} \delta_1'' + \delta_1'/r \\ \delta_2'' + \delta_2'/r \end{pmatrix} = M \begin{pmatrix} \delta_1 \\ \delta_2 \end{pmatrix}\tag{18}$$

$$\delta_3'' + \frac{\delta_3'}{r} = \left(\frac{1}{r^2} + \frac{\alpha^2}{2}(v_1^2 + v_2^2) \right) \delta_3,\tag{19}$$

where M is the mass matrix given by (9). So, for large r , we obtain $\delta_3 \propto K_1(m_Z r) \sim r^{-1/2} \exp(-m_Z r)$. Diagonalising the mass matrix M by writing $\begin{pmatrix} \delta_1 \\ \delta_2 \end{pmatrix} = \Delta_1 \vec{e}_1 + \Delta_2 \vec{e}_2$,

where $\vec{e}_1, \vec{e}_2$ are the eigenvectors of M , we get decoupled equations for Δ_1 and Δ_2 , yielding $\Delta_1 \propto K_0(m_{H_0}r)$ and $\Delta_2 \propto K_0(m_{h_0}r)$, demonstrating that δ_1 and δ_2 are linear combinations of $r^{-1/2} \exp(-m_{H_0}r)$ and $r^{-1/2} \exp(-m_{h_0}r)$. If m_{H_0} or m_{h_0} is greater than $2m_Z$, the asymptotics are slightly different^[14], with δ_3 source terms present in the Bessel equations for Δ_1, Δ_2 which cannot be neglected.

We note that for the particular value of the parameters

$$\tan \beta \equiv \frac{v_2}{v_1} = \left(\frac{\lambda_1 - \lambda_5}{\lambda_2 - \lambda_5} \right)^{\frac{1}{2}}, \quad (20)$$

setting $f = f_1/v_1 = f_2/v_2$ causes equations (14) and (15) to reduce to the same equation:

$$f'' + \frac{f'}{r} - \left(1 - \frac{\alpha}{2}Z\right)^2 \frac{f}{r^2} - ((\lambda_1 + \lambda_3)v_1^2 + (\lambda_3 + \lambda_5)v_2^2)(f^2 - 1)f = 0. \quad (21)$$

In this special case, $\begin{pmatrix} \delta_1 \\ \delta_2 \end{pmatrix} = \Delta_1 \begin{pmatrix} 1 \\ \tan \beta \end{pmatrix} + \Delta_2 \begin{pmatrix} \tan \beta \\ -1 \end{pmatrix}$. We know that $f_1/v_1 = f_2/v_2$, and hence $\delta_2/\delta_1 = \tan \beta$, yielding $\Delta_2 \equiv 0$. This shows that, despite the exponential dependence in Δ_2 decaying more slowly than that in Δ_1 , we cannot just ignore the more rapidly decaying Δ_1 when studying the asymptotic behaviour of δ_1 and δ_2 .

We shall examine this case in detail later since it leads to a considerable reduction in the complexity of the stability problem.

3. The Harmonic Decomposition of the Perturbations

The vortex solution given in (13) is not topologically stable. The Higgs vacuum is $S_3 \times S_3$ and the first homotopy group of this is trivial. We are investigating the metastability of the vortex solution i.e. whether it is a *local* maximum or minimum in configuration space. We consider infinitesimal perturbations of the vortex configuration and ask if the quadratic variation in the energy is positive or negative. We write

$$\begin{aligned} \Phi_1 &= \begin{pmatrix} \chi_1 \\ f_1(r)e^{i\theta} + \delta\phi_1 \end{pmatrix}, \quad \Phi_2 = \begin{pmatrix} \chi_2 \\ f_2(r)e^{i\theta} + \delta\phi_2 \end{pmatrix} \\ \vec{Z} &= -\frac{1}{r}Z(r)\hat{e}_\theta + \delta\vec{Z}, \quad \vec{A} = \vec{a}, \\ \vec{W}^{\bar{a}}\tau^{\bar{a}} &= \vec{W}_+\tau^- + \vec{W}_-\tau^+, \end{aligned} \quad (22)$$

where we find it convenient to use the complex notation

$$W_{\pm} = \frac{1}{\sqrt{2}} (W_1 \pm iW_2), \quad \tau^{\pm} = \frac{1}{\sqrt{2}} (\tau^1 \pm i\tau^2). \quad (23)$$

The perturbations can depend on the z -coordinate and the z -components of the vector fields can be non-zero also. However, since the vortex solution has translational invariance along the z -direction, it is easy to see that the z -dependence in the perturbations can be ignored and the z -components of the gauge fields can be set to zero. This follows from (4) where the relevant z -dependent terms in the integrand are:

$$\frac{1}{2}G_{i3}^a G_{i3}^a + \frac{1}{4}F_{i3}F_{i3} + (D_3\Phi_1)^{\dagger}(D_3\Phi_1) + (D_3\Phi_2)^{\dagger}(D_3\Phi_2) \quad (24)$$

This contribution to the energy is strictly non-negative and is made to vanish by setting the z -components of the gauge fields to zero and also considering the perturbations to be independent of the z -coordinate. For this reason, we shall drop all reference to the z -coordinate in the calculations below and it will be understood that the energy is actually the energy *per unit length* of the string.

The only contribution of the photon field is the second term of the integrand in (4) and this gives

$$\frac{1}{2}(\vec{\nabla} \times \vec{a})^2$$

which is positive and hence we set $\vec{a} = 0$.

The quadratic change in the energy may then be written as

$$\delta E = \int d^2x \left(\chi_{\bar{a}} \quad \phi_{\bar{a}}, \quad \delta \vec{Z}, \quad \vec{W}_- \right)^* \mathbf{O} \begin{pmatrix} \chi_{\bar{a}} \\ \phi_{\bar{a}} \\ \delta \vec{Z} \\ \vec{W}_- \end{pmatrix}, \quad (25)$$

where $\mathbf{O}$, the fluctuation operator, is a second order matrix differential operator. The perturbations will then decompose into channels labelled by the eigenvalues of the operators generating the continuous symmetries of $\mathbf{O}$ and by the eigenvalues of the discrete symmetries of $\mathbf{O}$. These symmetries are the subset of the symmetries of the energy density which are also symmetries of the vortex configuration (13).

It has a Z_2 symmetry under the action of the large $U(2)$ gauge transformation given by

$$\overline{U} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (26)$$

The perturbations of the string itself, $\delta\phi_{\bar{a}}$ and $\delta\vec{Z}$, lie in the $\overline{U} = 1$ channel and $\chi_{\bar{a}}$ and $\vec{W}^{\bar{a}}\tau^{\bar{a}}$ are in $\overline{U} = -1$. Hence these sets of perturbations decouple and we can write

$$\delta E = \delta E_1(\delta\phi_{\bar{a}}, \delta\vec{Z}) + \delta E_2(\chi_{\bar{a}}, W_- \tau_+). \quad (27)$$

The first term is positive since it corresponds to the stability problem of the vortex in the abelian Higgs case which is topologically stable as discussed in reference [15]. Hence we set $\delta\phi_{\bar{a}} = 0$, $\delta\vec{Z} = 0$.

The configuration is axially symmetric i.e. it is invariant under the action of the symmetry operator generated by the generalised angular momentum operator

$$K_z = L_z + S_z + I_z \quad (28)$$

where L_z and S_z are the usual orbital and spin pieces respectively of the spatial angular momentum operator given explicitly by

$$L_z = -i \frac{\partial}{\partial \theta} \mathbf{1}, \quad (S_z \vec{b})_j = -i \epsilon_{3jk} \vec{b}_k \mathbf{1}, \quad (29)$$

where $\mathbf{1}$ is the unit matrix and $\vec{b}$ is any vector field (note that S_z annihilates the Higgs doublets). I_z is composed of a $U(1)$ generator, Y and an $SU(2)$ generator, T^3

$$I_z = -\frac{1}{2}(Y - T^3). \quad (30)$$

We therefore perform a harmonic expansion of the perturbations,

$$\begin{aligned} \chi_{\bar{a}} &= \sum_m \chi_{\bar{a}}^m(r) e^{im\theta} \\ \vec{W}_- \tau^+ &= \sum_n \left(-iF_n(r) \hat{e}_r + \frac{\xi_n(r)}{r} \hat{e}_\theta \right) e^{i(n-1)\theta} \tau^+ \end{aligned} \quad (31)$$

where m and n take values from $-\infty$ to $+\infty$. It is clear that this is a decomposition into angular momentum modes, since

$$K_z(e^{im\theta}) = m e^{im\theta}, \quad K_z(e^{i(n-1)\theta} \tau^+) = n e^{i(n-1)\theta}. \quad (32)$$

The configuration is also CP-invariant i.e. it is invariant under the combination of reflection in the x-axis and complex conjugation. Therefore the channels may be labelled as CP-even and CP-odd. It is straightforward to check that the $U(2)$ gauge transformation $U = \text{diag}[i, 1]$ swaps the two channels and hence the two instability problems are exactly equivalent. Hence this supplies a reality condition and we may take $\chi_a^m, F_n(r)$ and $\xi_n(r)$ to be real.

4. Analysis of the Least Stable Mode

We have shown that the relevant energy variation can be decomposed into a sum over angular momentum modes

$$\delta E = \sum_m \delta E(\chi_a^m, F_m(r), \xi_m(r)). \quad (33)$$

Explicitly for the m^{th} mode we split the variation in the energy into three contributions

$$\delta E(\chi_a^m, F_m(r), \xi_m(r)) = E_h + E_c + E_W \quad (34)$$

where E_h arises solely from the perturbation in the Higgs fields, E_W is from the perturbation of the W-fields and E_c is from the interaction.

Then

$$\begin{aligned} E_h = 2\pi \int_0^\infty r dr \Big[& |\chi_1^{m'}|^2 + |\chi_2^{m'}|^2 + \frac{1}{r^2} \left(m + \frac{\alpha}{2} \cos 2\Theta_W Z \right)^2 (|\chi_1^m|^2 + |\chi_2^m|^2) \\ & + (\lambda_1 + \lambda_3)(f_1^2 - v_1^2) |\chi_1^m|^2 + (\lambda_2 + \lambda_3)(f_2^2 - v_2^2) |\chi_2^m|^2 \\ & + \lambda_3((f_1^2 - v_1^2) |\chi_2^m|^2 + (f_2^2 - v_2^2) |\chi_1^m|^2 + \lambda_4 |f_2 \chi_1^m - f_1 \chi_2^m|^2 \\ & + 2\lambda_5(f_1 f_2 - v_1 v_2) \chi_1^m \chi_2^m \Big] \end{aligned} \quad (35)$$

$$\begin{aligned} E_c = 2\pi g \int_0^\infty r dr \Big[& ((f_1' \chi_1^m - f_1 \chi_1^{m'}) + (f_2' \chi_2^m - f_2 \chi_2^{m'})) F_m \\ & - \frac{\xi_m}{r^2} (f_1 \chi_1 + f_2 \chi_2) (m + 1 - \alpha \sin^2 \Theta_W Z) \Big] \end{aligned} \quad (36)$$

$$E_W = \pi \int_0^\infty \frac{dr}{r} \left[2F_m \xi_m \gamma Z' + ((m - (1 - \gamma Z)) F_m - \xi_m')^2 + \frac{g^2(f_1^2 + f_2^2)}{2} (\xi_m^2 + r^2 F_m^2) \right] \quad (37)$$

An instability is most likely to develop in the core of the string, away from the vacuum manifold. Considering the Higgs perturbations, if we examine E_h (35) we see that a Higgs condensate in the core could lead to a decrease in the potential energy. The Higgs perturbations can only be non-vanishing in the core for $m=0$, else the field is singular there.

If we now consider E_W (37) we see that the only term that can make a negative contribution is the first term of the integrand. The physical interpretation of this instability has been elucidated by Perkins^[16] : the magnetic field $\vec{B} = \vec{\nabla} \times \vec{Z}$ in the core of the string couples to the anomalous magnetic moment of the W-bosons and they acquire a negative effective mass. The potential instability is thus a W-condensate in the core. If we perform the Taylor expansion around the origin

$$F_m = a_{m0} + a_{m2}r^2 + \dots, \quad \xi_m = rb_{m0} + r^3b_{m2} + \dots, \quad (38)$$

then we find that

$$\vec{W}_{-\tau^+} = \sum_n \begin{pmatrix} -i(a_{m0} \cos \theta - ib_{m0} \sin \theta) \\ -ia_{m0} \sin \theta + b_{m0} \cos \theta \end{pmatrix} e^{i(m-1)\theta} \tau^+ + O(r^2). \quad (39)$$

Then the requirement that the fields be non-singular at the origin gives that they can only be non-vanishing there for $m=0, 2$. First considering the case $m=2$, we then require $F_2 = \xi_2/r$, that is they have the same sign and the contribution is therefore positive since Z' is positive in the core. However, for the case $m=0$ the condition is that $F_0 = -\xi_0/r$ and the contribution is then negative.

We conclude that the least stable mode is the $m=0$ mode. This confirms the intuitive idea that we expect the least stable mode to have $K_z = 0$ since as $|K_z|$ increases one gets an increasing centrifugal barrier. We henceforth restrict ourselves to the case $m=0$. If we examine (35–37) we note that there are no F' terms (we drop the

m-script from now on). We therefore complete the squares and obtain

$$\begin{aligned}
\delta E = \pi \int_0^\infty \frac{dr}{r} & \left[\xi'^2 + \frac{1}{2} g^2 (f_1^2 + f_2^2) \xi^2 - \frac{Q^2}{P_+} - 2g(1 - \alpha \sin^2 \Theta_W Z) \xi (f_1 \chi_1 + f_2 \chi_2) \right. \\
& + 2r^2 (\chi_1'^2 + \chi_2'^2) + \frac{1}{2} (\alpha \cos 2\Theta_W Z)^2 (\chi_1^2 + \chi_2^2) \\
& + 2r^2 \left\{ (\lambda_1 + \lambda_3) (f_1^2 - v_1^2) \chi_1^2 + (\lambda_2 + \lambda_3) (f_2^2 - v_2^2) \chi_2^2 + \lambda_4 (f_2 \chi_1 - f_1 \chi_2)^2 \right. \\
& \left. \left. + \lambda_3 ((f_1^2 - v_1^2) \chi_2^2 + (f_2^2 - v_2^2) \chi_1^2) + 2\lambda_5 (f_1 f_2 - v_1 v_2) \chi_1 \chi_2 \right\} \right] \\
& + T(F, \chi_1, \chi_2, \xi)
\end{aligned} \tag{40}$$

where

$$T(F, \chi_1, \chi_2, \xi) = \pi \int_0^\infty \frac{dr}{r} \left(\sqrt{P_+} F + \frac{Q}{\sqrt{P_+}} \right)^2, \tag{41}$$

and

$$Q = \xi'(1 - \gamma Z) + \gamma Z' \xi - g r^2 (f_1 \chi_1' - f_1' \chi_1 + f_2 \chi_2' - f_2' \chi_2), \tag{42}$$

and

$$P_+ = (1 - \gamma Z)^2 + \frac{1}{2} g^2 r^2 (f_1^2 + f_2^2). \tag{43}$$

The contribution from $T(F, \chi_1, \chi_2, \xi)$ is positive and we have an algebraic equation for F , namely

$$F = -\frac{Q}{P_+}, \tag{44}$$

so that the contribution is made to vanish. We are now left with a problem in three variables and we express the change in the energy in the form

$$\delta E[\chi_1, \chi_2, \xi] = 2\pi \int dr r (\chi_1, \chi_2, \xi) \bar{\mathbf{O}} \begin{pmatrix} \chi_1 \\ \chi_2 \\ \xi \end{pmatrix}, \tag{45}$$

where $\bar{\mathbf{O}}$ is a 3×3 matrix differential operator.

We have not completely fixed the gauge, and hence there is a gauge zero mode which we must extract. Perturbations of the form

$$\delta \Phi = ig \psi \Phi_0, \quad \delta W_i^a \tau^a = -i D_{0i} \psi, \tag{46}$$

where ψ is a real $L(SU(2))$ valued function and the 0 subscript denotes the unperturbed fields, give an infinitesimal gauge transformation of the vortex solution (13). If we now

require that the gauge perturbations lie in (χ_1, χ_2, ξ) then we can only have

$$\psi = s(r) \begin{pmatrix} 0 & ie^{-i\theta} \\ -ie^{i\theta} & 0 \end{pmatrix} \quad (47)$$

where $s(r)$ is any smooth function. This means that perturbations given by

$$\begin{pmatrix} \chi_1 \\ \chi_2 \\ \xi \end{pmatrix} = s(r) \begin{pmatrix} -gf_1 \\ -gf_2 \\ 2(1 - \gamma Z) \end{pmatrix} \quad (48)$$

are pure gauge perturbations and are annihilated by $\bar{\mathbf{O}}$. Hence we need a basis for our physical perturbations orthogonal to this. We choose

$$\begin{aligned} \begin{pmatrix} \chi_1 \\ \chi_2 \\ v\xi \end{pmatrix} &= \begin{pmatrix} gf_1/2 \\ gf_2/2 \\ -v(1 - \gamma Z) \end{pmatrix} s_0(f_1^2 + f_2^2)^{-1/2} + \begin{pmatrix} f_2 \\ -f_1 \\ 0 \end{pmatrix} s_1(f_1^2 + f_2^2)^{-1/2} \\ &+ \begin{pmatrix} f_1(1 - \gamma Z) \\ f_2(1 - \gamma Z) \\ \frac{g}{2v}(f_1^2 + f_2^2) \end{pmatrix} \frac{s_2}{P}(f_1^2 + f_2^2)^{-1/2}, \end{aligned} \quad (49)$$

where v is a mass-scale so that the basis vectors have consistent dimensions and

$$P = (1 - \gamma Z)^2 + \frac{g^2}{4v^2}(f_1^2 + f_2^2). \quad (50)$$

Before substituting (49) into (45), to make the dependence on the parameters $t = \tan \beta = v_2/v_1$, $\sin^2 \Theta_W$, $\beta_i = \lambda_i/\alpha^2$ explicit we rescale the fields as follows

$$R = vr, \quad v_1 F_1 = f_1, \quad v_2 F_2 = f_2, \quad \bar{Z} = \frac{1}{2}\alpha Z, \quad (51)$$

where we now set

$$v = \alpha(v_1 v_2)^{\frac{1}{2}}. \quad (52)$$

P_+ and P are now re-expressed as

$$\begin{aligned} P_+ &= (1 - 2 \cos^2 \Theta_W \bar{Z})^2 + \frac{1}{2} R^2 \cos^2 \Theta_W F^2 \\ P &= (1 - 2 \cos^2 \Theta_W \bar{Z})^2 + \frac{1}{4} \cos^2 \Theta_W F^2 \end{aligned} \quad (53)$$

where

$$F^2 = \frac{1}{t} F_1^2 + t F_2^2. \quad (54)$$

On substitution of (49) into (45) it was a good check on our algebra that the s_0 terms vanished. After a lot of algebra we obtain

$$\delta E = \pi \int_0^\infty dR \left(\underline{s}'^T M \underline{s}' + \underline{s}^T N \underline{s}' + \underline{s}^T S \underline{s} \right) \quad (55)$$

where ' now denotes differentiation with respect to R , $\underline{s}^T = (s_1, s_2)$, and

$$M = \begin{pmatrix} M_{11} & 0 \\ 0 & M_{22} \end{pmatrix}, \quad N = \begin{pmatrix} 0 & N_{12} \\ 0 & 0 \end{pmatrix}, \quad S = \begin{pmatrix} S_{11} & S_{12} \\ S_{12} & S_{22} \end{pmatrix}. \quad (56)$$

We give the coefficients explicitly in the appendix. Note that M is diagonal as one would expect from choosing orthogonal coordinates. Also, it is of interest to note that N_{12} and S_{12} , the coefficients giving the coupling between s_1 and s_2 , are both proportional to $(F_1 F_2' - F_2 F_1')$ so that for the special choice of parameters (20), which gives $F_1 = F_2$, N_{12} and S_{12} are zero and the equations decouple.

We are only assured of a complete set of eigenvectors and hence of obtaining all possible eigenvalues when the problem is posed in a self-adjoint form. The problem as it is posed in (55) is not in self-adjoint form due to the non-vanishing coefficient N_{12} . Hence we must perform a further change of variables

$$\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} s_1 \\ s_2 + \lambda s_1 \end{pmatrix}. \quad (57)$$

If we then choose

$$\lambda' = \frac{N_{12}}{2M_{22}} = 2(1 - 2\cos^2 \Theta_W \bar{Z}) \frac{(F_1 F_2' - F_2 F_1')}{F^2},$$

which we solve together with the boundary condition $\lambda(0) = 0$ then the problem (55) is now expressed in self-adjoint form:

$$\delta E = \pi \int_0^\infty dR \left(\underline{u}'^T \bar{M} \underline{u}' + \underline{u}^T \bar{S} \underline{u} \right), \quad (58)$$

where trivially $\bar{M}$ and $\bar{S}$ are symmetric. Now to find the coupled Schrödinger equation we extremise δE subject to fixing the weighted norm of $\underline{u}$ via

$$2\pi \int_0^\infty R dR \underline{u}^T \bar{W} \underline{u} = 1,$$

where

$$\bar{W} = \begin{pmatrix} 1 + \lambda^2/P_+ & -\lambda/P_+ \\ -\lambda/P_+ & 1/P_+ \end{pmatrix}. \quad (59)$$

We then introduce the Lagrange multiplier μ and extremise

$$\pi \int_0^\infty dR (\underline{u}'^T \bar{M} \underline{u}' + \underline{u}^T \bar{S} \underline{u} - 2\mu R \underline{u}^T \bar{W} \underline{u}), \quad (60)$$

which gives the coupled Schrödinger equations

$$-(\bar{M} \underline{u}')' + \bar{S} \underline{u} = 2\mu R \bar{W} \underline{u}. \quad (61)$$

This is equivalent to extremising

$$\pi \int_0^\infty dR (\underline{s}'^T M \underline{s}' + \underline{s}^T N \underline{s}' + \underline{s}^T S \underline{s} - 2\mu R \underline{s}^T W \underline{s}) \quad (62)$$

where

$$W = \begin{pmatrix} 1 & 0 \\ 0 & 1/P_+ \end{pmatrix}, \quad (63)$$

which gives the coupled equations

$$\begin{aligned} 2(M_{11}s_1')' &= N_{12}s_2' + 2(S_{11} - 2\mu R)s_1 + 2S_{12}s_2 \\ 2(M_{22}s_2')' + (N_{12}s_1)' &= 2(S_{22} - 2\mu R/P_+)s_2 + 2S_{12}s_1. \end{aligned} \quad (64)$$

The boundary conditions on s_1 and s_2 are taken to be

$$s_1^2(0) + s_2^2(0) = 1, \quad s_1(\infty) = s_2(\infty) = 0, \quad (65)$$

which is permissible due to the linearity of the problem, and the ratio $s_1(0)/s_2(0)$ will vary according to the eigenvalue μ . It then follows that the stability of the string is dependent on μ , since for the solutions of (64),

$$\delta E = \mu \int_0^\infty R dR \underline{s}^T W \underline{s}, \quad (66)$$

and since W and $\bar{W}$ are strictly positive definite then if μ is negative the string is unstable, and if μ is positive the string is stable.

5. Numerical Studies

On making the change of variables (51), equations (14–16) become

$$F_1'' + \frac{F_1'}{R} - (1 - \bar{Z})^2 \frac{F_1}{R^2} - \frac{1}{t} (\beta_1 + \beta_3) (F_1^2 - 1) F_1 - t\beta_3 (F_2^2 - 1) F_1 - t\beta_5 (F_1 F_2 - 1) F_2 = 0 \quad (67)$$

$$F_2'' + \frac{F_2'}{R} - (1 - \bar{Z})^2 \frac{F_2}{R^2} - t(\beta_2 + \beta_3) (F_2^2 - 1) F_2 - \frac{1}{t} \beta_3 (F_1^2 - 1) F_2 - \frac{1}{t} \beta_5 (F_1 F_2 - 1) F_1 = 0 \quad (68)$$

$$\bar{Z}'' - \frac{\bar{Z}'}{R} + \frac{1}{2} (1 - \bar{Z}) F^2 = 0 \quad (69)$$

which are solved together with the boundary conditions

$$F_1(0) = F_2(0) = \bar{Z}(0) = 0, \quad F_1(\infty) = F_2(\infty) = \bar{Z}(\infty) = 1 \quad (70)$$

by relaxation. To determine the minimum eigenvalue μ , the equations (64) are then solved by relaxation on the same lattice. Since the couplings between s_1 and s_2 are small, the spectrum of eigenvalues can be split into two families; one in which $\underline{s}$ is mainly s_1 , and one in which $\underline{s}$ is mainly s_2 . These families are studied separately by solving equations (64) in two separate cases, namely by imposing either $s_1(0) = 1$ or $s_2(0) = 1$, to yield two minimal eigenvalues which we will call μ_1 and μ_2 respectively.

If we set $\tan \beta = 1$, $\beta_1 = \beta_2$, $\beta_3 = \beta_4 = \beta_5 = 0$, then condition (20) is satisfied and the s_1 and s_2 equations decouple. Writing $\Phi_1 = \Phi_2 = \Phi/\sqrt{2}$ causes our energy functional to reduce to that of the standard electroweak model, provided we set $\beta_1 = \beta_2 = \frac{1}{2}\beta$, where $\sqrt{\beta}$ is the ratio of the Higgs and Z masses. In this special case, we found the same region of stability for s_2 perturbations as that found in references [8,9], which provided a useful check on the reliability of our numerics.

It was found that μ_1 is typically positive, giving no unstable modes made principally of s_1 perturbations. Since μ_2 depends only very weakly on β_4 , a negative μ_1 can easily be made positive by increasing the value of β_4 slightly without seriously affecting μ_2 . In the search for parameters which give a stable vortex solution, the main difficulty is

in making μ_2 positive. Over a wide range of parameters, it was found that the crucial parameter in determining μ_2 is

$$\frac{m_{H_0}^2}{m_Z^2} = \frac{2t}{1+t^2} \left\{ c_1 + c_2 + \left(t + \frac{1}{t} \right) c_5 + \left[\left(c_1 - c_2 + \left(t - \frac{1}{t} \right) c_5 \right)^2 + c_3^2 \right]^{1/2} \right\}, \quad (71)$$

where $c_1 = (\beta_1 + \beta_3)/t$, $c_2 = t(\beta_2 + \beta_3)$, $c_3 = 2\beta_3 + \beta_5$, $c_5 = \beta_5/2$. Moving around the parameter space on curves of fixed m_{H_0}/m_Z typically results in little variation in μ_2 . Hence, to analyse the stability properties, we can restrict ourselves to the case where our energy functional reduces to that of the standard electroweak theory, thus reproducing the stability properties of references [8,9]. Numerical studies demonstrate that by tweaking the parameters in the model for fixed mass ratio (71) μ_2 can be slightly increased from the value in the electroweak model, but this increase is small. Some sample results are displayed in table 1. Hence, since the region of stability for strings in the standard electroweak model is a long way from experimentally acceptable values of the Weinberg angle and Higgs to Z mass ratio, the increase in the region of stability obtained by adding a second Higgs doublet is insufficient to lead to stable strings for realistic parameters.

Table 1

$\sin^2 \Theta_W$	$\tan \beta$	β_1	β_2	β_3	β_5	$\mathcal{R}$	μ_2
0.95	1.0	0.8	0.8	0.0	0.0	1.6	-0.12*
0.95	1.0	0.2	0.2	0.2	0.2	1.6	-0.12
0.95	1.7	1.17	0.8	-0.4	0.5	1.6	-0.10
0.95	2.8	0.44	0.35	-0.12	0.6	1.6	-0.13
0.95	1.0	1.6	0.0	0.0	0.0	1.6	-0.28
0.6	1.0	0.2	0.2	0.0	0.0	0.4	-0.57*
0.6	1.0	0.05	0.05	0.05	0.05	0.4	-0.57
0.6	0.8	0.04	0.17	0.02	0.05	0.4	-0.52
0.6	1.0	0.4	0.0	0.0	0.0	0.4	-0.52
0.6	0.2	0.208	0.0	0.0	0.0	0.4	-1.86
0.6	2.0	1.0	0.0	0.0	0.0	0.4	-0.45

In table 1, $\mathcal{R} = m_{H_0}^2/m_Z^2$, and $\beta_4 = 0$ in all cases. In the starred cases, our model reduces to the one doublet electroweak theory.

6. Conclusion

In section 2 we demonstrated that there are vortex solutions in the CP-invariant two Higgs doublet electroweak theory throughout the allowable parameter range. The ‘width’ of such vortices depends on the masses of the two CP-even neutral Higgs particles H^0 and h^0 , and on the Z -mass. In section 3 a general perturbation of such a vortex was decomposed into angular momentum eigenstates, and it was seen that the perturbations decouple into two parts – one the same as in the two singlet abelian Higgs model, and a second part arising from nonzero $\vec{W}$ and Higgs perturbations orthogonal to the original profiles. The CP-invariance of the model was used to decouple the real and imaginary parts of the perturbations. In section 4, the least stable mode in our expansion of the perturbations was picked out after linearisation, and a change of coordinates was performed to make explicit the component of this mode which was an infinitesimal gauge transformation. The change in energy was then expressed in self-adjoint form, to be analysed as a Sturm-Liouville eigenvalue problem. The numerical study of the perturbations was discussed in section 5, and it was discovered that the most important determining factor in the sign of the energy associated with the perturbation was the ratio of the masses of the H^0 and Z particles. The region of stability was only slightly larger than that for vortices in the one doublet electroweak model, and we conclude that for realistic values of the Weinberg angle and Higgs mass, vortex solutions in the two doublet electroweak model are unstable.

Acknowledgements

We thank Anne-Christine Davis for her advice and for encouraging us in doing this problem. Both authors acknowledge a SERC research student grant.

Appendix

In section 4. we showed that the stability problem is reduced to extremising

$$\pi \int_0^\infty dR \left(\underline{s}'^T M \underline{s}' + \underline{s}^T N \underline{s}' + \underline{s}^T S \underline{s} - 2\mu R \underline{s}^T W \underline{s} \right) \quad (72)$$

where ' denotes differentiation with respect to R .

$$M_{11} = P_+ M_{22} = 2R$$

$$N_{12} = 8R(1 - 2\cos^2 \Theta_W \bar{Z}) \frac{(F_1 F_2' - F_1' F_2)}{P_+ F^2}$$

$$\begin{aligned} S_{11} = & 2R \left[\frac{F_1 F_2' - F_1' F_2}{F^2} \right]^2 (1 - 2\cos^2 \Theta_W R^2 F^2 P_+^{-1}) + \frac{2}{R} \cos^2 \Theta_W \bar{Z}^2 + 2R\beta_4 F^2 \\ & + \frac{2R}{F^2} \left[(\beta_1 + \beta_3)(F_1^2 - 1)F_2^2 + (\beta_2 + \beta_3)(F_2^2 - 1)F_1^2 \right. \\ & \left. + \beta_3 \left(\frac{1}{t^2}(F_1^2 - 1)F_1^2 + t^2(F_2^2 - 1)F_2^2 \right) - 2\beta_5(F_1 F_2 - 1)F_1 F_2 \right] \end{aligned}$$

$$S_{12} = 4R \frac{(F_1 F_2' - F_1' F_2)}{P_+ F^2} \left[2\cos^2 \Theta_W \bar{Z}' - (1 - 2\cos^2 \Theta_W \bar{Z}) \frac{F'}{F} \right]$$

$$S_{22} = A_1 + A_2 - \frac{1}{2} A_3'$$

where

$$\begin{aligned} A_1 = & \frac{1}{8} \cos^4 \Theta_W \frac{R}{P^2 P_+} (FF')^2 - \frac{1}{2} \cos^2 \Theta_W \frac{P'}{RP^3} FF' + \frac{1}{4} \cos^2 \Theta_W \frac{P'^2}{RP^4} F^2 \\ & + \frac{1}{8} \cos^4 \Theta_W \frac{F^4}{RP^2} + 2\cos^2 \Theta_W (1 - 2\cos^2 \Theta_W \bar{Z})^2 \frac{\bar{Z}^2}{RP^2} \\ & - \cos^2 \Theta_W (1 - 2\cos^2 \Theta_W \bar{Z})(1 - 2\sin^2 \Theta_W \bar{Z}) \frac{F^2}{RP^2} \\ & + \frac{2R}{P^2} [2\cos^2 \Theta_W \bar{Z}' + (1 - 2\cos^2 \Theta_W \bar{Z})P'/P]^2 \\ & + \frac{2R}{P^2} (1 - 2\cos^2 \Theta_W \bar{Z})^2 \left[\left(\frac{F_1 F_2' - F_1' F_2}{F^2} \right)^2 - \frac{1}{2} \cos^2 \Theta_W \frac{F'^2}{P_+} (1 + R^2) \right] \\ & - \cos^2 \Theta_W \frac{F^2}{RP^2 P_+} \left[\left(\frac{1}{2} + R^2 \right) 2\cos^2 \Theta_W \bar{Z}' - \left(\frac{1}{2} - R^2 \right) (1 - 2\cos^2 \Theta_W \bar{Z}) P'/P \right]^2 \\ & - 2\cos^2 \Theta_W \frac{FF'}{RP^2 P_+} \left(\frac{1}{2} + R^2 \right) \times \left[\left(\frac{1}{2} + R^2 \right) 2\cos^2 \Theta_W \bar{Z}' (1 - 2\cos^2 \Theta_W \bar{Z}) \right. \\ & \left. - \left(\frac{1}{2} - R^2 \right) (1 - 2\cos^2 \Theta_W \bar{Z})^2 P'/P \right] \end{aligned}$$

$$\begin{aligned}
A_2 = & \frac{2R}{P^2 F^2} (1 - 2 \cos^2 \Theta_W \bar{Z})^2 \left[(\beta_1 + \beta_3)(F_1^2 - 1) \frac{F_1^2}{t^2} + (\beta_2 + \beta_3)(F_2^2 - 1) F_2^2 t^2 \right. \\
& \left. + \beta_3 ((F_1^2 - 1) F_2^2 + (F_2^2 - 1) F_1^2) + 2\beta_5 (F_1 F_2 - 1) F_1 F_2 \right] \\
A_3 = & \frac{1}{R P^2 P_+} [\cos^2 \Theta_W R^2 F F' - 2(1 - 2 \cos^2 \Theta_W \bar{Z}) 2 \cos^2 \Theta_W \bar{Z}'] \times \\
& \left[2R^2 (1 - 2 \cos^2 \Theta_W \bar{Z})^2 + \frac{1}{4} \cos^2 \Theta_W F^2 \right] - 4R \frac{P'}{P P_+}
\end{aligned}$$

References

- [1] T. VACHASPATI, M. BARRIOLA *Phys. Rev. Lett.* **69** (1992), 1867
M. BARRIOLA, T. VACHASPATI, M. BUCHER preprint TUTP-93-7
- [2] T. VACHASPATI, A. ACHÚCARRO *Phys. Rev.* **D44** (1991), 3067
- [3] H. B. NIELSEN, P. OLESEN *Nucl. Phys.* **B61** (1973), 45
- [4] M. HINDMARSH *Phys. Rev. Lett.* **68** (1992), 1263
- [5] A. ACHÚCARRO, K. KUIJKEN, L. PERIVOLAROPOULOS, T. VACHASPATI, *Nucl. Phys.* **B388** (1992), 435
- [6] T. VACHASPATI *Phys. Rev. Lett.* **68** (1992), 1977
- [7] Y. NAMBU *Nucl. Phys.* **B130** (1977), 505
- [8] M. JAMES, T. VACHASPATI, L. PERIVOLAROPOULOS *Phys. Rev.* **D46** (1992), 5232
- [9] M. JAMES, T. VACHASPATI, L. PERIVOLAROPOULOS *Nucl. Phys.* **B395** (1993), 534
- [10] R. HOLMAN, S. HSU, T. VACHASPATI, R. WATKINS *Phys. Rev.* **D46** (1993), 5352
- [11] R. BRANDENBERGER, A-C. DAVIS *Phys. Lett.* **B308** (1993), 79
- [12] T. VACHASPATI, R. WATKINS preprint hep-ph/921184
- [13] J.F. GUNION, H.E. HABER, G.L. KANE, S. DAWSON ‘The Higgs Hunter’s Guide’
Addison Wesley (1989)
- [14] L. PERIVOLAROPOULOS preprint (1993)
- [15] H.S. LA preprint hep-ph/9302220
- [16] W.B. PERKINS *Phys. Rev.* **D47** (1993), 5224